


**FIGURE 3.** Predicted protein structure of the double mutant E746V-L747P. This structural rearrangement allows ATP (orange and blue chemical structure) binding, even in the presence of gefitinib (purple chemical structure), in a favorable position for hydrolysis. Visual analysis of the protein structures based on native EGFR alone, in complex either with tyrosine kinase inhibitor or ATP analogue, mutated EGFR with gefitinib from Protein Data Bank (<http://www.rcsb.org/pdb/home/home.do>. Accessed February 10, 2015; PDB: 4WRG, 4WKQ, 4HJO, 4G5J, 4I23, 3VJO, 4I22), was carried over using the program COOT and figure was drawn using the program Chimera.



**Simona Coco, PhD**  
**Anna Truini, MS**  
**Irene Vanni, MS**  
**Carlo Genova, MD**

Lung Cancer Unit  
 IRCCS AOU San Martino-IST  
 Istituto Nazionale per la Ricerca sul  
 Cancro  
 Genova, Italy

**Camillo Rosano, PhD**  
 Biopolymers and Proteomics Unit  
 IRCCS AOU San Martino-IST  
 Istituto Nazionale per la Ricerca sul  
 Cancro  
 Genova, Italy

**Maria Giovanna Dal Bello, PhD**  
**Angela Alama, PhD**  
 Lung Cancer Unit  
 IRCCS AOU San Martino-IST  
 Istituto Nazionale per la Ricerca sul  
 Cancro  
 Genova, Italy

**Roberta Venè, PhD**  
 Molecular Oncology and  
 Angiogenesis Unit  
 IRCCS AOU San Martino-IST  
 Istituto Nazionale per la Ricerca sul  
 Cancro  
 Genova, Italy

**Erika Rijavec, MD**  
**Giulia Barletta, MD**  
**Federica Biello, MD**  
 Lung Cancer Unit  
 IRCCS AOU San Martino-IST

Istituto Nazionale per la Ricerca sul  
 Cancro  
 Genova, Italy

**Francesco Boccardo, MD**  
 Dipartimento di Medicina Interna e  
 Specialità Mediche (DIMI)  
 IRCCS AOU San Martino-IST  
 Istituto Nazionale per la Ricerca sul  
 Cancro  
 Genova, Italy  
 UOC Clinica di Oncologia Medica,  
 IRCCS AOU San Martino-IST  
 Istituto Nazionale per la Ricerca sul  
 Cancro  
 Genova, Italy

**Francesco Grossi, MD**  
 Lung Cancer Unit  
 IRCCS AOU San Martino-IST  
 Istituto Nazionale per la Ricerca sul  
 Cancro  
 Genova, Italy

## ACKNOWLEDGEMENTS

*The present study was supported by "Fondazione Carige" (2013.0277-12) and the Italian Ministry of Health (GR 2011-12; 02350922; SC). We thank Drs. Cristina Bruzzo and Zita Cavalieri for technical support. We are grateful to the tissue bank Centro di Risorse Biologiche-IST (CRB-IST).*

## REFERENCES

1. van der Wekken AJ, Stigt JA, A't Hart N. A novel EGFR mutation in exon 19 showed

stable disease after TKI treatment. *J Thorac Oncol* 2012;7:e8.

2. Wu JY, Yu CJ, Chang YC, Yang CH, Shih JY, Yang PC. Effectiveness of tyrosine kinase inhibitors on "uncommon" epidermal growth factor receptor mutations of unknown clinical significance in non-small cell lung cancer. *Clin Cancer Res* 2011;17:3812-3821.
3. Niho S, Kubota K, Goto K, et al. First-line single agent treatment with gefitinib in patients with advanced non-small-cell lung cancer: a phase II study. *J Clin Oncol* 2006;24:64-69.
4. Gajiwala KS, Feng J, Ferre R, et al. Insights into the aberrant activity of mutant EGFR kinase domain and drug recognition. *Structure* 2013;21:209-219.
5. He M, Capelletti M, Nafa K, et al. EGFR exon 19 insertions: a new family of sensitizing EGFR mutations in lung adenocarcinoma. *Clin Cancer Res* 2012;18:1790-1797.

## MicroRNA Assays for Diagnosis Lung Cancer Biopsy

### To the Editor:

The recent report on "MicroRNA Assays to Distinguish Squamous Cell Carcinoma from Adenocarcinoma in Lung Cancer Biopsies" is very interesting.<sup>1</sup> Patnaik et al.<sup>1</sup> noted that "histotypic microRNA assays can aid the subtyping of non-small-cell lung cancer biopsies as

Address for correspondence: Viroj Wiwanitkit, MD, Surin Rajabhat University, Surin, Thailand. E-mail: [wviroj@yahoo.com](mailto:wviroj@yahoo.com)

DOI: 10.1097/JTO.0000000000000528

Copyright © 2015 by the International Association for the Study of Lung Cancer  
 ISSN: 1556-0864/15/1006-0e52

adenocarcinoma (AC) or squamous cell carcinoma (SCC) by standard histopathologic methods.” In fact, adding a new diagnostic tool to the classical diagnostic tool might improve the diagnostic ability. Focusing on the use of microRNA assays, there are still left concern and questions on the cost effectiveness, availability, and complexity of the tests. These points have to be further discussed. Focusing on MiR-205 MicroRNA, its diagnostic value for differentiating between AC and SCC is still controversial. Some previous reports showed limitation of its ability to diagnose SCC.<sup>2</sup> Also, the MiR-205 can also increase in the case with severe inflammation and benign tumor.<sup>3</sup> The possibility of false-positive because of noncancerous lesion has to be further studied.

**Viroj Wiwanitkit, MD**  
Public Health Curriculum  
Surin Rajabhat University  
Surin, Thailand

#### REFERENCES

1. Patnaik S, Mallick R, Kannisto E, et al. MiR-205 and MiR-375 MicroRNA assays to distinguish squamous cell carcinoma from adenocarcinoma in lung cancer biopsies. *J Thorac Oncol* 2015;10:446–453.
2. Manikandan M, Deva Magendhra Rao AK, Munirajan AK. Altered levels of miR-21, miR-125b-2\*, miR-134, miR-155, miR-184, and miR-205 in oral squamous cell carcinoma and association with clinicopathological characteristics. *J Oral Pathol Med*. 2014 December 8 [epub ahead of print].
3. Nurul-Syakima AM, Learn-Han L, Yoke-Kqueen C. miR-205 in situ expression and localization in head and neck tumors—a tissue array study. *Asian Pac J Cancer Prev* 2014;15:9071–9075.

## Reply to “MiR-205 and miR-375 microRNA Assays to Distinguish Squamous Cell Carcinoma From Adenocarcinoma in Lung Cancer Biopsies”

#### In Response:

We thank Dr. Wiwanitkit for his comments on our study.<sup>1</sup> The microRNA-based assay described in it

requires the quantification of only four RNAs (*miR-21*, *miR-205*, and *miR-375*, and *RNU6B*). As already noted by us in the publication, this can be conveniently done in any laboratory with a quantitative polymerase chain reaction machine, with time and material costs similar to those for immunohistochemistry-based diagnosis of non-small-cell lung cancer histology. However, the suitability of the assay for biospecimens with less than 90% tumor content has not been assessed by us. In our study, microdissection of tumor-containing regions of biopsied material was performed for 76% of biopsies to have ≥90% tumor content in the specimens that were used for RNA extraction for the microRNA-based assay.

The studies on the association of *miR-205* with severe inflammation and benign tumor and the lack of a differential expression of this microRNA between normal and tumor tissues, which Dr. Wiwanitkit refers to, concern oral cancer and not cancer of the lung. In case of the latter, a significantly higher expression of *miR-205* in lung squamous cell carcinoma tissue compared with normal lung, or lung tissue with adenocarcinoma or benign diseases has been noted by many<sup>2–5</sup> and in Figure 1 of our article.<sup>1</sup>

**Santosh K. Patnaik, MBBS, PhD**  
**Sai Yendamuri, MD**  
Department of Thoracic Surgery  
Roswell Park Cancer Institute  
Buffalo, New York

**Reema Mallick, MD**  
Department of Surgery  
University of Minnesota  
Minneapolis, Minnesota

**Samjot S. Dhillon, MD**  
Department of Medicine  
(Pulmonary/Thoracic Oncology)  
Roswell Park Cancer Institute  
Buffalo, New York

Address for correspondence: Samjot S. Dhillon, Department of Medicine (Thoracic Oncology and Pulmonary Medicine), Roswell Park Cancer Institute, Elm and Carlton Streets, Buffalo, New York. E-mail: samjot.dhillon@roswellpark.org  
DOI: 10.1097/JTO.0000000000000539  
Copyright © 2015 by the International Association for the Study of Lung Cancer  
ISSN: 1556-0864/15/1006-0e53

#### REFERENCES

1. Patnaik S, Mallick R, Kannisto E, et al. MiR-205 and MiR-375 MicroRNA Assays to Distinguish Squamous Cell Carcinoma from Adenocarcinoma in Lung Cancer Biopsies. *J Thorac Oncol* 2015;10:446–453.
2. Lebanony D, Benjamin H, Gilad S, et al. Diagnostic assay based on hsa-miR-205 expression distinguishes squamous from non-squamous non-small-cell lung carcinoma. *J Clin Oncol* 2009;27:2030–2037.
3. Zhang YK, Zhu WY, He JY, et al. miRNAs expression profiling to distinguish lung squamous-cell carcinoma from adenocarcinoma subtypes. *J Cancer Res Clin Oncol* 2012;138:1641–1650.
4. Bishop JA, Benjamin H, Cholak H, Chajut A, Clark DP, Westra WH. Accurate classification of non-small cell lung carcinoma using a novel microRNA-based approach. *Clin Cancer Res* 2010;16:610–619.
5. Jiang M, Zhang P, Hu G, et al. Relative expressions of miR-205-5p, miR-205-3p, and miR-21 in tissues and serum of non-small cell lung cancer patients. *Mol Cell Biochem* 2013;383:67–75.

## Reply to “Better Prognostic Models May Result in Improved Patient Selection for Adjuvant Therapies After Complete Resection of Solitary Fibrous Tumors of the Pleura”

#### In Response:

We would like to thank Dr. Tapias and Dr. Lanuti for their comments on our recent article reporting on a multicenter cohort of 68 patients with solitary fibrous tumors of the pleura (SFTP), who were analyzed for the complete course of the disease in a routine practice setting.<sup>1</sup> We acknowledge that our recurrence rate of 30%

Address for correspondence: Olivier Bylicki, MD, Service de Pneumologie, Respiratory Medicine and Thoracic Oncology Service, Hôpital Louis Pradel, 28 avenue doyen Lépine, 69677 Lyon (Bron) Cedex, France. E-mail: bylicki.olivier@yahoo.fr  
DOI: 10.1097/JTO.0000000000000557  
Copyright © 2015 by the International Association for the Study of Lung Cancer  
ISSN: 1556-0864/15/1006-0e53